

臺灣綜合大學系統 108 學年度學士班轉學生聯合招生考試試題

科目名稱	線性代數	類組代碼	A07.C11
		科目碼	A0702
※本項考試依簡章規定各考科均「不可以」使用計算機			本科試題共計 3 頁
一、單選題（占 28 分）請於答案卷上作答，否則不予計分			
說明：第 1 題至第 4 題，每題有 8 個選項，答案以英文字母大寫 ABCDEFGH 作答。各題答對者，得 7 分。答錯、未作答或多於一個答案者，該題以零分計算。			
<p>1. (7pts) <math>A</math> is a <math>m</math>-by-<math>n</math> matrix, and the system <math>A\vec{x} = \vec{0}</math> has a unique solution <math>\vec{x} = \vec{0}</math>.</p> <ul style="list-style-type: none"> <li>(i) <math>\forall \vec{b} \in \mathbb{R}^m</math>, there exists at least one solution to <math>A\vec{x} = \vec{b}</math>.</li> <li>(ii) If there is a solution to <math>A\vec{x} = \vec{b}</math>, then this is a unique solution.</li> <li>(iii) <math>m \leq n</math>.</li> </ul> <p>Which of the above statements are correct (if any)? _____.</p> <p>A. (i); B. (ii); C. (iii); D. (i)(ii); E. (i)(iii); F. (ii)(iii); G. (i)(ii)(iii); H. None.</p>			
<p>2. (7pts) <math>A</math> and <math>B</math> are two <math>n</math>-by-<math>n</math> real matrices.</p> <ul style="list-style-type: none"> <li>(i) <math>\det(AB) = \det(A)\det(B)</math>.</li> <li>(ii) <math>AB</math> and <math>BA</math> have the same eigenvalues.</li> <li>(iii) <math>I</math> is the <math>n</math>-by-<math>n</math> identity matrix. If <math>AB = I</math>, then <math>BA = I</math>.</li> </ul> <p>Which of the above statements are correct (if any)? _____.</p> <p>A. (i); B. (ii); C. (iii); D. (i)(ii); E. (i)(iii); F. (ii)(iii); G. (i)(ii)(iii); H. None.</p>			
<p>3. (7pts) <math>A</math> and <math>B</math> are two similar <math>n</math>-by-<math>n</math> real matrices.</p> <ul style="list-style-type: none"> <li>(i) <math>A</math> and <math>B</math> have the same characteristic polynomial.</li> <li>(ii) <math>A</math> and <math>B</math> have the same eigenvectors.</li> <li>(iii) If <math>A</math> is symmetry, then <math>B</math> is symmetry.</li> </ul> <p>Which of the above statements are correct (if any)? _____.</p> <p>A. (i); B. (ii); C. (iii); D. (i)(ii); E. (i)(iii); F. (ii)(iii); G. (i)(ii)(iii); H. None.</p>			
<p>4. (7pts) Matrix <math>A</math> is symmetric positive definite and matrix <math>Q</math> is orthogonal.</p> <ul style="list-style-type: none"> <li>(i) <math>Q^T A Q</math> is a diagonal matrix.</li> <li>(ii) <math>Q^T A Q</math> is symmetric positive definite.</li> <li>(iii) All pivots of <math>Q^T A Q</math> (without row changes) are positive.</li> </ul> <p>Which of the above statements are correct (if any)? _____.</p> <p>A. (i); B. (ii); C. (iii); D. (i)(ii); E. (i)(iii); F. (ii)(iii); G. (i)(ii)(iii); H. None.</p>			

臺灣綜合大學系統 108 學年度學士班轉學生聯合招生考試試題

科目名稱	線性代數	類組代碼	A07.C11
		科目碼	A0702
※本項考試依簡章規定各考科均「不可以」使用計算機			本科試題共計 3 頁
二、單選題（占 72 分）請於答案卷上作答，否則不予計分			
說明：1. 第 5 題至第 14 題，每題有 9 個選項，答案以英文字母大寫 ABCDEFGHI 作答。 2. 第 5 至 12 題答對給 7 分，第 13 至 14 題答對給 8 分。答錯、未作答或多於一個答案者，該題以零分計算。			
5. (7pts) Matrix $A = \begin{bmatrix} 1/2 & -4 & -2 \\ 0 & 2 & 2 \\ 0 & 0 & -1 \end{bmatrix}$ . $A^{-1} = \begin{bmatrix} * & a & b \\ * & * & c \\ * & * & * \end{bmatrix}$ . Then $a + b + c = \underline{\hspace{2cm}}$ . A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
6. (7pts) Linear transformation $T: \mathbf{R}^2 \rightarrow \mathbf{R}^3$ satisfies $T \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ a \end{bmatrix}$ and $T \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ b \\ 6 \end{bmatrix}$ . If $T$ is <u>not</u> one-to-one, then $a + b = \underline{\hspace{2cm}}$ . A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
7. (7pts) Matrix $A = \begin{bmatrix} 0 & 2 & 4 & 2 \\ 1 & 12 & 5 & 4 \\ 2 & 22 & 6 & 6 \\ 3 & 32 & 7 & 8 \\ 4 & 42 & 8 & 0 \end{bmatrix}$ . $\text{Rank}(A) = \underline{\hspace{2cm}}$ . A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
8. (7pts) $T \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{bmatrix} -1 & -1 & 3 & 0 & 3 \\ 0 & -1 & 2 & 1 & 1 \\ 1 & -2 & 3 & 3 & 0 \end{bmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix}$ is a linear operator from $\mathbf{R}^5$ to $\mathbf{R}^3$ . $\text{Ker}(T) = \left\{ \alpha \begin{pmatrix} a \\ d \\ 1 \\ 0 \\ 0 \end{pmatrix} + \beta \begin{pmatrix} b \\ e \\ 0 \\ 1 \\ 0 \end{pmatrix} + \gamma \begin{pmatrix} c \\ f \\ 0 \\ 0 \\ 1 \end{pmatrix}, \forall \alpha, \beta, \gamma \in \mathbf{R} \right\}$ . Then $a + c + e = \underline{\hspace{2cm}}$ . A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
9. (7pts) Matrix $A = \begin{bmatrix} 3 & 0 & 5 \\ 2 & -1 & 2 \\ 1 & 0 & 2 \end{bmatrix}$ , $A^{-1} = aA^2 + bA + cI$ . Then $a + b + c = \underline{\hspace{2cm}}$ . A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			

臺灣綜合大學系統 108 學年度學士班轉學生聯合招生考試試題

科目名稱	線性代數	類組代碼	A07.C11
		科目碼	A0702
※本項考試依簡章規定各考科均「不可以」使用計算機			本科試題共計 3 頁
10. (7pts) Given five points $(x_1, y_1) = (-2, 9)$ , $(x_2, y_2) = (-1, 2)$ , $(x_3, y_3) = (0, -6)$ , $(x_4, y_4) = (1, 0)$ , $(x_5, y_5) = (2, 0)$ . The constants $C, D, E$ are undetermined real numbers. Find the best fitting parabola $Cx^2 + Dx + E$ to the above five points, using least squares. In other words, the best solution $C, D, E$ is the one that minimizes $\sum_{i=1}^5 (Cx_i^2 + Dx_i + E - y_i)^2$ .			
Then $C - D - E = \underline{\hspace{2cm}}$ .			
A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
11. (7pts) Let $x, y \in \mathbb{R}$ . The minimum of $\frac{5x^2 - 2xy + 5y^2}{x^2 + y^2}$ is $a$ . Then $a = \underline{\hspace{2cm}}$ .			
A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
12. (7pts) $P_3$ is the set of all real polynomials of degree less or equal to 3 with ordered basis $\beta_P = \{x^3, x^2, x, 1\}$ . $M_{2 \times 2}$ is the set of all real 2-by-2 matrices with ordered basis $\beta_M = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$ .			
A linear operator $T: P_3 \rightarrow M_{2 \times 2}$ is defined as $T(f(x)) = \begin{bmatrix} f(-1) & f(0) \\ f(1) & f(2) \end{bmatrix}$ .			
$[T]_{\beta_P}^{\beta_M} = \begin{bmatrix} a & * & * & * \\ 0 & 0 & 0 & b \\ * & * & c & * \\ * & d & * & * \end{bmatrix}$ is the matrix representation of $T$ with respect to $\beta_P$ and $\beta_M$ .			
Then $a + b + c + d = \underline{\hspace{2cm}}$ .			
A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
13. (8pts) Matrix $A = \begin{bmatrix} 9/2 & 7/2 \\ 7/2 & 9/2 \end{bmatrix}$ , $A^{1/3} = \begin{bmatrix} a & c \\ b & d \end{bmatrix}$ . Then $a + b + c + d = \underline{\hspace{2cm}}$ .			
A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			
14. (8pts) $P_2$ is the set of all real polynomials of degree less or equal to 2. Define the inner product on $P_2$ by $\langle f, g \rangle = \int_0^1 f(x)g(x)dx$ , $\forall f, g \in P_2$ . Let $S$ be the subspace of $P_2$ spanned by $\{x^2, x\}$ . For $u(x) = 1$ , there exists $v \in S$ and $w \in S^\perp$ so that $u = v + w$ . Find $w(x)$ in the form $w(x) = 1 + ax + bx^2$ . Then $a + 3b = \underline{\hspace{2cm}}$ .			
A. 1; B. 2; C. 3; D. 4; E. 5; F. 6; G. 7; H. 8; I. 9.			